FOCUSED PROFESSIONAL REVIEW / 1 OF 2

Urology: evaluate the whole pathway.

A focused reading of rehabilitation and erectile-function research.

16 / Tenuto 2 / Conference abstract / 2025

Penile rehabilitation after radical prostatectomy

Uncontrolled combined intervention | 16 participants · 6 months

Mean IIEF erectile-function score was 22 before surgery and 19 six months after surgery, indicating partial recovery. 15 of 16 patients found the device easy to use; 13 reported improved intimacy and communication with their partners.

Uncontrolled combined treatment with tadalafil and a penile pump. The device contribution cannot be isolated. The postoperative score was lower than the preoperative score; 11 is the lower end of the postoperative range, not a baseline mean.

Journal of Sexual Medicine 22, Supplement 2

Read source publication

For the clinical discussion

Identify the population, the model, accompanying treatment and the relevant endpoint. Do not promise independent device efficacy from a combined intervention. Confirm follow-up, current instructions and the clinician responsible for reassessment.

Educational research summary, not a treatment protocol or an additional product indication. Review the full source, exact model instructions and individual suitability. This review edition requires clinical approval before wider distribution.

Read the full evidence library

Model guides and support

FOCUSED PROFESSIONAL REVIEW / 2 OF 2

Evidence, with the right context.

19 / Tenuto Mini / Journal correspondence / 2025

A vibrating constriction ring for psychogenic erectile dysfunction

Uncontrolled preliminary study | 30 participants · 8 weeks (company summary)

Mean IIEF-5 score increased from 14.44 to 22.31 (approximately 55%), according to MV Health's summary. Mean treatment satisfaction was reported as 78% on the EDITS measure.

Uncontrolled preliminary research, published as correspondence. Full numerical results require access to the complete article.

International Journal of Impotence Research · online 2025; volume 38, 2026

Read source publication

For the clinical discussion

Identify the population, the model, accompanying treatment and the relevant endpoint. Do not promise independent device efficacy from a combined intervention. Confirm follow-up, current instructions and the clinician responsible for reassessment.

Educational research summary, not a treatment protocol or an additional product indication. Review the full source, exact model instructions and individual suitability. This review edition requires clinical approval before wider distribution.

Read the full evidence library

Model guides and support